

学校编码：10384

分类号

密级

学号：22620061152373

UDC _____

厦 门 大 学

硕 士 学 位 论 文

假俭草草坪抗旱节水及抗寒护绿
的化学调控技术研究

The study of water-saving and green-maintaining technique
by means of chemical regulation for
Eremochloa ophiuroides turf management

李 雄

指导教师姓名：胡宏友 讲师

专业名称：环境工程

论文提交日期：2009年7月

论文答辩日期：2009年8月

2009年9月

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下，独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果，均在文中以适当方式明确标明，并符合法律规范和《厦门大学研究生学术活动规范（试行）》。

另外，该学位论文为()课题(组)的研究成果，获得()课题(组)经费或实验室的资助，在()实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称，未有此项声明内容的，可以不作特别声明。)

声明人（签名）：

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文,并向主管部门或其指定机构送交学位论文(包括纸质版和电子版),允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索,将学位论文的标题和摘要汇编出版,采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于:

() 1.经厦门大学保密委员会审查核定的保密学位论文,
于 年 月 日解密,解密后适用上述授权。

() 2.不保密,适用上述授权。

(请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文,未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的,默认为公开学位论文,均适用上述授权。)

声明人(签名):

年 月 日

摘 要

草坪草普遍蒸腾耗水量大,且暖季型草坪草不耐冬季低温,枯黄期长,这是草坪养护管理中存在的两个主要问题。本研究以园林上广泛应用的暖季型草坪草假俭草为材料,采用盆栽和野外实验相结合筛选高效抗旱及促越冬期生长的配方,为城市草坪草养护提供配套的节水、节能及提高景观质量的管理技术。

在假俭草抗旱节水技术的研究上,本文首先探讨了干旱胁迫下假俭草各生长生理指标动态变化。结果表明,干旱处理 10 d 后,未喷施药剂的假俭草叶绿素含量急剧下降,叶片游离脯氨酸含量和可溶性糖含量迅速增加,膜脂过氧化产物 MDA 大量积累,开始呈现干旱胁迫临界期生理变化症状,而喷施外源药剂(水杨酸、CPPU)可使假俭草干旱胁迫的临界期由 10 d 延长至 13-15 d,即浇水时间间隔延长 30-50%,这对于草坪草的节水灌溉十分有利;在此基础上,研究进一步选择壳聚糖、多效唑、水杨酸、CPPU、黄腐酸等 5 种外源药剂进行抗旱性比较试验,结果表明,在所选择的浓度范围内,壳聚糖处理组生物量最高,并具有最大的根冠比,且叶片游离脯氨酸和丙二醛(MDA)含量也处于较低水平,这表明壳聚糖提高假俭草抗旱性能要优于其他 4 种药剂。此外,通过抗旱机制的初步研究表明,壳聚糖是通过增强蛋白质的稳定性,提高抗氧化酶活性,减缓丙二醛的积累以及提高叶绿素含量等等途径来达到提高假俭草抗旱性的目的,其中,250 mg·L⁻¹壳聚糖处理下的假俭草叶绿素含量及生长生物量较其他处理组高、MDA 含量则相对最低,均接近或达到正常浇水处理组的水平。综合分析表明,对于暖季型假俭草草坪,按 400 ml·m⁻²喷施 250 mg·L⁻¹壳聚糖溶液,可以使草坪正常浇水间隔时间延长 30-50%,能显著发挥抗旱节水效果。

在假俭草抗寒护绿技术研究上,利用 L₉(3⁴) 正交试验设计方法,本文设计了植物生长调节剂、速效肥与有机肥 3 因子 3 水平正交混施试验处理越冬期假俭草。与清水对照相比,正交混施处理组假俭草叶面积指数、叶绿素含量、叶片游离脯氨酸和可溶性糖含量均显著提高,同时根部根系活力、叶片抗氧化酶系(SOD、POD、CAT)活性也显著增加,叶片丙二醛含量则显著降低。处理组整体上促进了假俭草越冬期生长及抗冷性能的提高,显著改善了其冬季景观效果。

因子间极差分析表明,主效因子随生长和抗性指标的不同而存在差异,表明混施时各因子在保持假俭草草坪冬季良好生长、提高抗性生理功能上所起的作用具有互补性。综合各指标正交分析的结果,提出了冬季提高假俭草抗寒性的最优混施配方,即多效唑、三十烷醇、CPPU、尿素和壳聚糖分别按 300、2、20、1500 和 1000 mg·L⁻¹配制成液肥按 200 ml·m⁻²叶面喷施。

关键词：假俭草 抗旱 抗寒 节水 护绿 化学调控

Abstract

Turf often consumes large quantity of water through its strong leaf transpiration. Meanwhile, warm-season turfgrass is subjected to cold-stress, wither and turn yellow during winter. In order to solve these problems, this research focuses on exploring the corresponding techniques to reduce the water-consuming as well as to improve the overwintering landscape quality of urban gardening turf. *Eremochloa ophiuroides* (Munro.) Hack, one of the main warm-season turfgrass species has been now widely used in the urban gardening landscape, was selected as the study material for this research. Pot and field experiments were conducted to screen the corresponding chemical formula to improve the drought-resistance and cold-resistance of the warm-season turfgrass.

The water-saving technique for *E. ophiuroides* turf management was explored through a set of drought-stress experiments. Firstly, the results of the experiment for the study of dynamic response of *E. ophiuroides* to drought showed that the physiological drought-stress indexes of the grass without exogenous reagents application would change 10 days after simulated drought stress, for example, the chlorophyll content began to decrease significantly, and free proline and soluble sugar content began to increase rapidly, meanwhile, malondialdehyde (MDA) was also determined to accumulate in great quantity. However, for those treatments applied with exogenous reagents, the changes were observed 13-15 days after the same drought stress condition. It suggested that, the watering interval time could be increased by 30-50% through applying corresponding exogenous reagents, this is quite beneficial to water-saving irrigation of turfgrass. Secondly, five exogenous reagents, which are chitosan, paclobutrazol, salicylic acid, CPPU, fulvic acid were selected for a screening test, the results showed that, among the five reagents, the value of biomass and root-shoot ratio of *E. ophiuroides* treated with chitosan were higher, meanwhile, the contents of free proline and MDA were lower than those treated by other reagents. It suggested that, compared with the other four kinds of exogenous reagents, chitosan was better to improve drought-resistance of *E. ophiuroides*. Further study revealed the chlorophyll content and biomass of *E. ophiuroides* in 250 mg·L⁻¹ chitosan treatment group was higher than those in 500 and

1000 mg·L⁻¹ chitosan treatment groups as well as the control, however, it was reversed for the MDA content. No significant differences of all the growth and physiological indexes were founded between the 250 mg·L⁻¹ chitosan treatment and the normal watering treatment. It suggested that chitosan can improve the drought-resistant ability of *E. ophiuroides* by enhancing its protein stability, improving the activities of antioxidant enzymes (SOD, POD and CAT), reducing the accumulation of MDA. Overall, foliar application of 250 mg·L⁻¹ chitosan solution with 400 ml·m⁻² can improve the drought-resistance of *E. ophiuroides* significantly. Under the drought condition, the interval time of turf irrigation could be increased by 30-50% in comparasion with the control.

In the study of promoting growth and prolonging green-maintaining period of warm-season turfgrass, a field experiment was carried out to treat *E. ophiuroides* with plant growth regulators, fast-released fertilizers and organic fertilizers by means of L₉(3⁴) orthogonal test. Compared with the control, the levels of LAI, chlorophyll, free proline and soluble sugar in the leaves of the treatment groups significantly improved, as well as the root activities, the activities of antioxidant enzymes, however, the contents of MDA decreased significantly. The growth, cold-resistance and the landscape quality of the treatment groups significantly promoted. The results of range analysis showed that the main effective factor varied with test indexes, which meant that plant growth regulators and fertilizers are complementary to promote the growth and enhance the physiological functions of turfgrass during winter. To improve the integrated quality of *E. ophiuroides* during winter, an optimum formula was proposed through comprehensively analyzing the growth and cold-resistant indexes, that is: a liquid fertilizer, containing paclobutrazol, triacontanol, CPPU, urea and chitosan with their concentrations as 300, 2, 20, 1500, and 1000 mg·L⁻¹ respectively, should be applied to the turf by foliage spray with 200 ml·m⁻².

Keywords: *Elemochloa ophiuroides*; Drought resistance; Cold resistance; Water-saving; Green-maintaining; Chemical regulation

目 录

中文摘要	I
英文摘要	III
表索引	XI
图索引	XIII
第一章 文献综述	1
1.1 引言	1
1.2 草坪草抗旱技术进展	2
1.2.1 选育抗旱性品种	2
1.2.2 抗旱锻炼	2
1.2.3 合理施肥	2
1.2.4 外源物质的调节	3
1.2.5 真菌、细菌的应用	3
1.2.6 其他管理手段	4
1.2.7 小结	4
1.3 暖季型草坪草抗寒技术进展	5
1.3.1 选育抗寒性品种	5
1.3.2 抗寒锻炼	6
1.3.3 合理施肥	6
1.3.4 植物生长调节剂的应用	8
1.3.5 其他管理手段	9
1.3.6 小结	10
1.4 总结	11
第二章 材料与方法	12
2.1 研究的思路与框架	12
2.2 研究地点概况	13
2.3 假俭草及其草坪的生物学属性	13
2.4 供试草皮的盆栽	13
2.5 假俭草抗旱节水技术的盆栽试验设计	14

2.5.1 外源抗旱药剂的选择.....	14
2.5.2 假俭草对干旱胁迫的响应特性的盆栽试验.....	15
2.5.3 外源药剂抗旱效果盆栽筛选试验.....	15
2.5.4 壳聚糖抗旱剂施用最佳浓度试验.....	16
2.6 假俭草越冬期护绿的化学调控技术试验	17
2.7 指标测定	18
2.7.1 生长及生物量.....	18
2.7.2 根系活性.....	18
2.7.3 叶绿素含量.....	18
2.7.4 可溶性糖含量.....	19
2.7.5 游离脯氨酸含量.....	19
2.7.6 清除活性氧酶系活性、丙二醛 (MDA) 和可溶性蛋白含量 ...	19
2.8 草坪综合质量评价方法	20
2.9 数据统计和处理	21
第三章 假俭草抗旱节水剂的筛选及其作用机制的研究	22
3.1 外源药剂处理下假俭草对干旱胁迫的响应特性	22
3.1.1 叶绿素含量动态.....	22
3.1.2 可溶性糖、游离脯氨酸含量动态.....	23
3.1.3 丙二醛及可溶性蛋白含量动态.....	24
3.1.4 清除活性氧酶系活性动态.....	25
3.1.5 讨论.....	27
3.1.6 小结.....	29
3.2 5种外源药剂提高假俭草抗旱能力的效果比较	29
3.2.1 对生物量的影响比较.....	30
3.2.2 对根冠比的影响比较.....	31
3.2.3 对丙二醛 (MDA) 含量的影响比较	31
3.2.4 对游离脯氨酸含量的影响比较.....	31
3.2.5 讨论.....	32
3.2.6 小结.....	35
3.3 壳聚糖抗旱剂施用浓度优化及其抗旱作用机制初步研究.....	35
3.3.1 光合色素及生长指标.....	35
3.3.2 清除活性氧酶系活性.....	39

3.3.3 丙二醛 (MDA) 含量	41
3.3.4 抗旱性生理指标	42
3.3.5 盆栽土的含水率	44
3.3.6 讨论	44
3.3.7 小结	47
3.4 总结	48
第四章 假俭草抗寒护绿技术研究	50
4.1 植物生长调节剂和肥料混施对假俭草越冬期生长和生理的影响	50
4.1.1 生长及生物量	50
4.1.2 根系活力及叶绿素含量	51
4.1.3 清除活性氧酶系活性	52
4.1.4 可溶性糖和可溶性蛋白含量	53
4.1.5 游离脯氨酸和丙二醛 (MDA) 含量	54
4.1.6 草坪草综合质量	55
4.2 讨论	56
4.2.1 调节剂和肥料混施对假俭草冬季营养生长的影响	56
4.2.2 调节剂和肥料混施对假俭草冬季活性氧代谢的影响	57
4.2.3 调节剂和肥料混施对假俭草抗寒性物质含量的影响	58
4.2.4 草坪综合质量评价体系的合理性及配方优化分析	59
4.3 总结	60
第五章 结论与展望	61
5.1 结论	61
5.2 存在问题及展望	62
5.2.1 存在问题	62
5.2.2 展望	63
参 考 文 献	64
附录 研究生期间发表的论文、参与的课题及其他成果	73
致 谢	74

CONTENT

Abstract in Chinese	I
Abstract in English	III
Table index	XII
Figure index	XIV
Chapter 1 Review	1
1.1 Introduction	1
1.2 Research progress on drought resistance technique of turfgrass	2
1.2.1 Breeding of drought-resistance varieties	2
1.2.2 Drought hardening	2
1.2.3 Fertilization	2
1.2.4 Exogenous reagents regulation	3
1.2.5 Application of fungi and bacteria.....	3
1.2.6 Other management means.....	4
1.2.7 Conclusion	4
1.3 Reseach progress on cold hardiness technique of warm-season turfgrass	5
1.3.1 Breeding of cold hardiness varieties	5
1.3.2 Cold hardening	6
1.3.3 Fertilization	6
1.3.4 Plant growth regulators application	8
1.3.5 Other management means.....	9
1.3.6 Conclusion	10
1.4 Summarization	11
Chapter 2 Materials and methods	12
2.1 Research thinking and framework	12
2.2 General situation of experimental field	13
2.3 Introduction of <i>Eremochloa ophiuroides</i> (Munro.) Hack	13
2.4 Potted culture of <i>Eremochloa ophiuroides</i> (Munro.) Hack	13
2.5 Pot experiment design of drought resistance and water saving technique .	14
2.5.1 Exogenous reagents selection	14
2.5.2 Response characteristics of <i>Eremochloa ophiuroides</i> (Munro.) Hack to drought stress	15
2.5.3 Screening exogenous reagents for their drought resistance effect.....	15

2.5.4 Optimum concentration of chitosan as a drought resistant reagent	16
2.6 Chemical regulation experiment on improving the cold-resistant and green-maintaining ability of <i>Eremochloa ophiuroides</i> (Munro.) Hack during winter	17
2.7 Index measurement	18
2.7.1 Growth and biomass	18
2.7.2 Root activity	18
2.7.3 Chlorophyll content	18
2.7.4 Soluble sugar content	19
2.7.5 Free proline content	19
2.7.6 Activities of reactive-oxygen-scavenging enzymes, MDA and soluble protein contents	19
2.8 Integrated quality evaluation	20
2.9 Data statistics and analysis	21
Chapter 3 The Study on screening turf drought-resistant reagents and its acting mechanism	22
3.1 Danymic response of <i>Eremochloa ophiuroides</i> (Munro.) Hack to drought with two exogeous reagents.....	22
3.1.1 Danymic of chlorophyll content	22
3.1.2 Danymics of proline and soluble sugar contents	23
3.1.3 Danymics of MDA and soluble protein contents	24
3.1.4 Danymics of reactive-oxygen-scavenging enzyme activities	25
3.1.5 Discussion	27
3.1.6 Conclusion	29
3.2 Comparison of five exogenous reagents on improving drought resistance of <i>Eremochloa ophiuroides</i> (Munro.) Hack.....	29
3.2.1 Comparison of biomass.....	30
3.2.2 Comparison of R/C	31
3.2.3 Comparison of MDA content	31
3.2.4 Comparison of free proline content	31
3.2.5 Discussion	32
3.2.6 Conclusion	35
3.3 Concentration optimazation of chitosan application and primary study on its acting mechanism	35
3.3.1 Photosynthetic pigment and growth.....	35
3.3.2 Activities of reactive-oxygen-scavenging enzymes	39
3.3.3 MDA content.....	41

3.3.4 Drought resistance indices	42
3.3.5 Soil water content	44
3.3.6 Discussion	44
3.3.7 Conclusion	47
3.4 Summarization.....	48
Chapter 4 Study of cold resistant and green-maintaining technique of <i>Eremochloa ophiuroides</i> (Munro.) Hack during winter	50
4.2 Effects of mixed applying PGRs and fertilizers on growth and cold-resistance of <i>Eremochloa ophiuroides</i> (Munro.) Hack during winter	50
4.1.1 Growth and biomass	50
4.1.2 Root activity and chlorophyll content.....	51
4.1.3 Activities of reactive-oxygen-scavenging enzymes.....	52
4.1.4 Soluble sugar and soluble protein content	53
4.1.5 Free proline and MDA content	54
4.1.6 Integrated quality evaluation.....	55
4.2 Discussion	56
4.2.1 Effects of mixed applying PGRs and fertilizers on growth of <i>Eremochloa ophiuroides</i> (Munro.) Hack during winter.....	56
4.2.2 Effects of mixed applying PGRs and fertilizers on active oxygen metabolism of <i>Eremochloa ophiuroides</i> (Munro.) Hack during winter.....	57
4.2.3 Effects of mixed applying PGRs and fertilizers on substances related with cold resistance of <i>Eremochloa ophiuroides</i> (Munro.) Hack during winter.....	58
4.2.4 Rationality of integrated quality evaluation and formula optimization	59
4.3 Conclusion	60
Chapter 5 Conclusion and prospect	61
5.1 Conclusion.....	61
5.2 Problems existed and prospect	62
5.2.1 Problems existed	62
5.2.2 Prospect.....	63
Reference	64
Appendix	73
Acknowledgement	74

表索引

表 1 抗旱试验处理药剂及浓度.....	16
表 2 $L_9(3^4)$ 因子水平表.....	18
表 3 生长调节剂及肥料混施对假俭草生长、生物量影响的多重比较及极差分析	50
表 4 生长调节剂及肥料混施影响假俭草根系活力和叶绿素含量的多重比较及极差分析	51
表 5 生长调节剂及肥料混施影响假俭草叶片活性氧清除酶系活性的多重比较及极差分析.....	52
表 6 生长调节剂及肥料混施影响假俭草可溶性糖和可溶性蛋白含量的多重比较及极差分析.....	53
表 7 生长调节剂及肥料混施影响假俭草 MDA 及游离脯氨酸含量的多重比较及极差分析	54
表 8 生长调节剂及肥料混施对假俭草综合质量的多重比较及极差分析	56

Table index

Table 1 Exogenous reagents selected and their concentrations for drought resistance experiments	16
Table 2 Factors and levels of $L_9(3^4)$ orthogonal treatment	18
Table 3 Multiple comparison and range analysis of the growth and biomass of <i>Eremochloa ophiuroides</i> (Munro.) Hack among the treated groups with plant growth regulators and fertilizers	50
Table 4 Multiple comparison and range analysis of the root activities and chlorophyll contents of <i>Eremochloa ophiuroides</i> (Munro.) Hack among the treated groups with plant growth regulators and fertilizers	51
Table 5 Multiple comparison and range analysis of the activities of active oxygen scavenging enzymes of <i>Eremochloa ophiuroides</i> (Munro.) Hack among the treated groups with plant growth regulators and fertilizers	52
Table 6 Multiple comparison and range analysis of soluble sugar and soluble protein contents of <i>Eremochloa ophiuroides</i> (Munro.) Hack among the treated groups with plant growth regulators and fertilizers.....	53
Table 7 Multiple comparison and range analysis of MDA and free proline contents of <i>Eremochloa ophiuroides</i> (Munro.) Hack among the treated groups with plant growth regulators and fertilizers.....	54
Table 8 Multiple comparison and range analysis of integrated quality of <i>Eremochloa ophiuroides</i> (Munro.) Hack among the treated groups with plant growth regulators and fertilizers	56

图索引

图 1 假俭草抗旱节水及抗寒护绿研究技术路线.....	12
图 2 持续干旱下 CPPU 和水杨酸处理对假俭草叶绿素含量动态的影响	22
图 3 持续干旱下 CPPU 和水杨酸处理对假俭草可溶性糖和游离脯氨酸含量动态的影响	24
图 4 持续干旱下 CPPU 和水杨酸处理对假俭草可溶性蛋白和丙二醛 (MDA) 含量动态的影响.....	25
图 5 持续干旱下 CPPU 和水杨酸处理对假俭草清除活性氧酶系活性动态的影响	26
图 6 干旱胁迫下 5 种药剂处理对假俭草的生物量和根冠比的影响	30
图 7 干旱胁迫下 5 种药剂处理对假俭草丙二醛和游离脯氨酸含量的影响	32
图 8 壳聚糖对干旱胁迫下假俭草叶绿素含量的影响 (与正常浇水处理组比较)	36
图 9 壳聚糖对干旱胁迫下假俭草草层高度的影响 (与正常浇水处理组比较)	37
图 10 壳聚糖对干旱胁迫下假俭草生物量的影响 (与正常浇水处理组比较)	38
图 11 壳聚糖对干旱胁迫下假俭草根冠比的影响 (与正常浇水处理组比较)	39
图 12 壳聚糖对干旱胁迫下假俭草活性氧清除酶系活性的影响 (与正常浇水处理组比较)	40
图 13 壳聚糖对干旱胁迫下假俭草丙二醛 (MDA) 含量的影响 (与正常浇水处理组比较)	42
图 14 壳聚糖对干旱胁迫下假俭草可溶性蛋白、脯氨酸和可溶性糖含量的影响 (与正常浇水处理组比较)	43
图 15 壳聚糖对干旱胁迫下假俭草土壤含水率的影响 (与正常浇水处理组比较)	44
图 16 处理组与对照组综合质量比较.....	55

Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

厦门大学博硕士论文摘要库